



CAMCEVI[®]
(leuprolide) 42 mg
injection emulsion



BE PREPARED

LEARN ABOUT TREATING YOUR
ADVANCED PROSTATE CANCER

IMPORTANT SAFETY INFORMATION

What is CAMCEVI[®]?

CAMCEVI is an androgen deprivation therapy (ADT) used for treatment of adult patients with advanced prostate cancer.

You should not take CAMCEVI if you have ever had an allergic reaction to any of the ingredients in CAMCEVI or to any other medications called gonadotropin-releasing hormone (GnRH) agonists.

Fully open brochure for the complete Important Safety Information.



CAMCEVI[®]
(leuprolide) 42 mg
injection emulsion



UNDERSTANDING YOUR PLAN

Whether you have been living with advanced prostate cancer for a while or are newly diagnosed, it's important to know that there are treatment options. Your healthcare professional will work with you on your treatment plan, which may now include CAMCEVI[®].

WHAT IS GnRH AGONIST HORMONE-BLOCKING THERAPY?

GnRH stands for *gonadotropin-releasing hormone*. It's a type of medicine that blocks the testicles from making sex hormones.¹

WHY LOWER YOUR TESTOSTERONE?

Testosterone is the male sex hormone that is made in the testicles. It "feeds" prostate cancer, which allows it to grow. Hormone therapy will stop your body from making testosterone or it will block what testosterone does in the body.²

IMPORTANT SAFETY INFORMATION

CAMCEVI may:


- Cause an increase in your testosterone level during the initial few weeks of treatment, which may cause temporary worsening of symptoms or additional signs and symptoms of prostate cancer. You may experience a temporary increase in bone pain. You may have trouble with urination or develop abnormal amount of pressure on your spinal cord which can result in paralysis, which may be life-threatening.
- Cause your blood sugar to increase, including increasing your risk of developing diabetes. You should have your blood sugar monitored while taking CAMCEVI.
- Increase your risk of heart problems including heart attack, sudden death, or stroke. It can also cause your heart to beat irregularly.
- Cause some patients to have seizures.

Fully open brochure for the complete Important Safety Information.

HOW IT WORKS

1


TESTOSTERONE FEEDS CANCER CELLS



Most of a man's testosterone is made in the testicles. Testosterone encourages the growth of prostate cancer cells.²

2

CAMCEVI® BLOCKS TESTOSTERONE PRODUCTION AT THE SOURCE



CAMCEVI prevents the testicles from making testosterone. This makes it harder for the tumor to grow.²

3

TESTOSTERONE LEVELS DROP³



In about 4 weeks, your testosterone levels are expected to drop to very low levels.³ Your healthcare professional will monitor your testosterone levels to know how well CAMCEVI is working.

NOTE: After a first injection of CAMCEVI, testosterone levels may go up for a short while.³ This is known as a *flare*, and it is common. Your healthcare professional will monitor you throughout treatment to know how well CAMCEVI is working. Once testosterone levels are reduced, studies show they stay low for 6 months.³ Then it's time for another CAMCEVI treatment.

UNDERSTANDING SIDE EFFECTS

Side effects can be common with hormone-blocking therapies like CAMCEVI. It's important to know what the common side effects of CAMCEVI are, so you can talk to your healthcare professional about them. **Call your healthcare professional for medical advice about side effects.**

IMPORTANT SAFETY INFORMATION

CAMCEVI may:

- Cause fetal harm if administered to a pregnant woman.
- Cause impotence and impair fertility in males of reproductive potential.

Fully open brochure for the complete Important Safety Information.

accord BioPharma

CAMCEVI, Accord BioPharma, and associated logos are trademarks of Accord BioPharma. All other trademarks are property of respective owners. ©2022 Accord BioPharma. All rights reserved. This content is intended for U.S. audiences only. PA-CAM-1015 – 7/22


CAMCEVI®
(leuprolide) 42 mg
injection emulsion

THE MOST COMMON SIDE EFFECTS:

- Hot flash³
- High blood pressure³
- Reaction where you received your shot, which may cause temporary burning/stinging, pain, bruising, or redness³
- Upper respiratory tract infections³
- Muscle and bone pain³
- Tiredness³
- Pain in extremity, such as the neck, shoulders, arms, and legs³

IMPORTANT SAFETY INFORMATION

What is CAMCEVI®?

CAMCEVI is an androgen deprivation therapy (ADT) used for treatment of adult patients with advanced prostate cancer.

You should not take CAMCEVI if you have ever had an allergic reaction to any of the ingredients in CAMCEVI or to any other medications called gonadotropin-releasing hormone (GnRH) agonists.

CAMCEVI may:

- Cause an increase in your testosterone level during the initial few weeks of treatment, which may cause temporary worsening of symptoms or additional signs and symptoms of prostate cancer. You may experience a temporary increase in bone pain. You may have trouble with urination or develop abnormal amount of pressure on your spinal cord which can result in paralysis, which may be life-threatening.
- Cause your blood sugar to increase, including increasing your risk of developing diabetes. You should have your blood sugar monitored while taking CAMCEVI.
- Increase your risk of heart problems including heart attack, sudden death, or stroke. It can also cause your heart to beat irregularly.
- Cause some patients to have seizures.
- Cause fetal harm if administered to a pregnant woman.
- Cause impotence and impair fertility in males of reproductive potential.

The most common side effects of CAMCEVI include hot flush, high blood pressure, reaction where you received your shot which may cause temporary burning/stinging, pain, bruising or redness, upper respiratory tract infections, muscle and bone pain, tiredness, and pain in extremity. Tell your healthcare provider if you experience a rash or severe injection site reaction.

For more information, talk to your healthcare provider.

These are not all the possible side effects of CAMCEVI. For additional information, see the full prescribing information or ask your healthcare professional or pharmacist.

You are encouraged to report negative side effects of prescription drugs to the FDA at 1-800-FDA-1088 or visit www.fda.gov/medwatch.

CHOOSING YOUR TREATMENT

Your healthcare professional has several options for managing advanced prostate cancer. There are many reasons why CAMCEVI® might be right for you. CAMCEVI is administered by your healthcare provider once every 6 months.



Leuprolides have shown to be effective in treating advanced prostate cancer³



Reduction in testosterone levels by week 4 after the initial dose³



Consistent reduction of testosterone levels from week 4 through week 48. By week 4, your testosterone levels are expected to be low enough to not promote cancer growth³




Subcutaneous injection means the injection is given in the fatty tissue, just under the skin³

RECEIVING YOUR INJECTION

You can only receive CAMCEVI injections from your healthcare provider. The proper dose is injected into the fatty tissue just under your skin in your abdomen.³ As long as CAMCEVI is working for you, your healthcare provider will repeat the injection every 6 months.

Injection Sites



**LEARN WHAT TO EXPECT
DURING YOUR TREATMENT.**



Visit [CAMCEVI.com](https://www.camcevi.com)

Please see accompanying full Prescribing Information.

RESOURCES FOR YOU

MORE WAYS TO CARE

The **AccordCares™** program offers support and resources to you, your loved ones, and your healthcare professional, including financial assistance and more. Services include copay assistance, alternate coverage programs, and assistance to your healthcare provider to ensure eligible coverage for your treatment is verified.



Patients who meet eligibility criteria may be eligible for payment assistance.

Download your enrollment form at **accordcaresportal.com**


CONTACT ACCORDCARES

Phone: (866) 258-7151 | **Address:** PO Box 221643, Charlotte, NC, 28222
Fax: (855) 558-6304 | **Hours:** Mon-Fri, 8am - 8pm EST

GET THE FULL STORY AT STORYMD

Learn more about the symptoms, tests, and treatments for prostate cancer, presented as informative visuals that can be discussed with your healthcare provider.

+STORYMD



Scan code to get started.

References:

1. National Cancer Institute. NCI Dictionaries. GnRH Agonist. Accessed July 14, 2022. <https://www.cancer.gov/publications/dictionaries/cancer-terms/def/gnrh-agonist>
2. NCCN Guidelines for Patients. Prostate Cancer Advanced Stage, 2020. Published 2020. Accessed July 14, 2022. <https://www.nccn.org/patients/guidelines/content/PDF/prostate-advanced-patient.pdf>
3. CAMCEVI. Prescribing information. Accord BioPharma; May 2021.